

Duality mappings within three-dimensional nonlinear electrodynamics

Mauricio Cataldo*

*Departamento de Física, Facultad de Ciencias,
Universidad del Bío-Bío, Avenida Collao 1202,
Casilla 5-C, Concepción, Chile.*

(Dated: February 1, 2008)

Abstract: In three-dimensional Einstein-Maxwell gravity the electrostatic Bañados-Teitelboim-Zanelli solution and the magnetostatic Hirschmann-Welch solution are connected by a duality mapping. Here we point out that a similar duality mapping exists among circularly symmetric electrostatic and magnetostatic spacetimes, and electric and magnetic stationary solutions, for a nonlinear electrodynamics coupled to three-dimensional Einstein gravity.

PACS numbers: 04.20.Jb

I. INTRODUCTION

In three-dimensional Einstein-Maxwell gravity exist the electric and magnetic analogs to the Reissner-Nordström-Kottler solution. They are the Bañados-Teitelboim-Zanelli (BTZ) black hole [1]

$$ds^2 = -(r^2/l^2 - M - q_e^2 \ln r) d\tilde{t}^2 + \frac{dr^2}{(r^2/l^2 - M - q_e^2 \ln r)} + r^2 d\tilde{\theta}^2, \quad (1)$$

where the radial electric field is given by $E = q_e/r$, and the Hirschmann-Welch (HW) magnetic solution [2]

$$ds^2 = -(r^2/l^2 - M) d\tilde{t}^2 + \frac{r^2 dr^2}{(r^2/l^2 - M)(r^2 + q_m^2 \ln(r^2/l^2 - M))} + (r^2 + q_m^2 \ln(r^2/l^2 - M)) d\tilde{\theta}^2, \quad (2)$$

where the pseudoscalar magnetic field is given by $B = q_m/r$. Here $\Lambda = -1/l^2 < 0$, q_e is the electric charge, q_m magnetic charge and M is the mass of the BTZ black hole. Both solutions are asymptotically anti-de Sitter like the Reissner-Nordström-Kottler solution for $\Lambda < 0$.

In [3] it is shown that there exist a duality mapping which connects the BTZ black hole and the HW solution. This mapping is given by

$$q_e \longrightarrow i q_m, \quad \tilde{t} \longrightarrow i \theta, \quad \tilde{\theta} \longrightarrow i t.$$

In this case the BTZ black hole takes the form

$$ds^2 = -r^2 d\tilde{t}^2 + \frac{dr^2}{(r^2/l^2 - M + q_m^2 \ln r)} + (r^2/l^2 - M + q_m^2 \ln r) d\tilde{\theta}^2. \quad (3)$$

It could be seen, by introducing a new coordinate r , that the metric (3) becomes (2).

This property is related to the fact that in (2+1)-dimensions the Maxwell tensor has only three independent components, two for the vector electric field and one for the scalar field. This means that the duality transformation which takes place for the Reissner-Nordström-Kottler black hole do not take place now. The reason is that in (3+1)-dimensions the Maxwell tensor and its dual are two-forms and in (2+1)-dimensions the Maxwell tensor is a two-form and its dual is a one form $*F_a = 1/2 \epsilon_{abc} F^{bc}$. This implies that the electric and magnetic counterparts are quite different.

In this letter we point out that a similar duality mapping exist among circularly symmetric electrostatic and magnetostatic spacetimes for a nonlinear electrodynamics coupled to three-dimensional Einstein gravity.

The letter is organized as follows. The next section deals with duality mappings among electrostatic and magnetostatic Einstein-nonlinear Maxwell spacetimes and we discuss some specific examples of nonlinear magnetic solutions. In Sec. III we discuss stationary Einstein-nonlinear Maxwell three-dimensional spacetimes. The electric and magnetic spinning cases are discussed separately. Finally, in Sec. IV some properties of these spinning nonlinear three-dimensional spacetimes are discussed.

II. DUALITY MAPPINGS WITHIN STATIC EINSTEIN-NONLINEAR ELECTRODYNAMICS

In (2+1)-dimensions, electromagnetic theories can be constructed from Lagrangians (without higher derivatives) depending on a single (non-vanishing) invariant $F = \frac{1}{4} F_{ab} F^{ab}$, which can be expressed in terms of the electric (vector) and magnetic (pseudoscalar) fields: in a lorentzian frame, for an observer moving with the 3-velocity v^a , the electric and the magnetic fields are cor-

*mcataldo@ubiobio.cl

respondingly defined as

$$E_a = F_{ab}v^b, \quad B = \frac{1}{2}\epsilon_{abc}F_{bc}v^a,$$

where latin indices run the values 0, 1, 2 and ϵ_{abc} is the totally anti-symmetric Levi-Civita symbol with $\epsilon_{012} = 1$, usually the v^a is oriented along the time coordinate, i.e, $v^a = \delta_t^a$, with such a choice

$$E_a = F_{a0}, \quad B = F_{12}.$$

Thus the invariant can be expressed by

$$F \equiv \frac{1}{4}F^{ab}F_{ab} = \frac{1}{2}(B^2 - E^2).$$

In general one can construct a (2+1)-Einstein theory coupled to a nonlinear electrodynamics starting from the action

$$S = \int \sqrt{-g} \left(\frac{1}{16\pi}(R - 2\Lambda) + L(F) \right) d^3x,$$

with the electromagnetic Lagrangian $L(F)$ unspecified explicitly at this stage; physically one requires the Lagrangian to coincide with the Maxwell one at small values of the electromagnetic fields, $L(F)_{Maxwell} = -F/4\pi$. We are using units in which $c = G = 1$, because of the ambiguity in the definition of the gravitational constant (there is not newtonian gravitational limit in (2+1)-dimensions) we prefer to maintain the factor $1/16\pi$ in the action to keep the parallelism with (3+1)-gravity. Varying this action with respect to gravitational field gives the Einstein equations

$$G_{ab} + \Lambda g_{ab} = 8\pi T_{ab},$$

where

$$T_{ab} = g_{ab}L(F) - F_{ac}F_b{}^c L_{,F}, \quad (4)$$

while the variation with respect to the electromagnetic potential A_a entering in $F_{ab} = A_{b,a} - A_{a,b}$, yields the electromagnetic field equations

$$\nabla_a (F^{ab} L_{,F}) = 0, \quad (5)$$

where $L_{,F}$ denotes the derivative of $L(F)$ with respect to F .

Concrete solutions to the dynamical equations we shall seek in the form

$$ds^2 = -e^{2\alpha(r)}dt^2 + e^{2\beta(r)}dr^2 + e^{2\gamma(r)}d\phi^2, \quad (6)$$

where α , β and γ are unknown functions of the variable r . We have chosen this form because if the structural functions are given by $e^{2\alpha(r)} = e^{-2\beta(r)}$ and $e^{2\gamma(r)} = r^2$ a magnetic field can not be obtained [4].

The general form of the electromagnetic field tensor which shares the static circularly symmetric spacetime is given by

$$F_M = E_r dr \wedge dt + B dr \wedge d\theta. \quad (7)$$

This gives us the following invariant F (from now on $E_r \equiv E$):

$$F = \frac{1}{4}F_{ab}F^{ab} = \frac{1}{2}(-E^2 e^{-2\alpha-2\beta} + B^2 e^{-2\beta-2\gamma}). \quad (8)$$

Thus from (4) we have for nonvanishing components

$$\begin{aligned} T_{00} &= -e^{2\alpha}L(F) - E^2 e^{-2\beta} L_{,F} \\ T_{11} &= e^{2\beta}L(F) + (E^2 e^{-2\alpha} - B^2 e^{-2\gamma}) L_{,F} \\ T_{22} &= e^{2\gamma}L(F) - B^2 e^{-2\beta} L_{,F} \\ T_{02} &= -EB e^{-2\beta} L_{,F}. \end{aligned} \quad (9)$$

Then the Einstein equations are

$$\begin{aligned} e^{-2\beta}(\gamma'\beta' - \gamma'' - \gamma'^2) = \\ \Lambda + \kappa [-L(F) - E^2 e^{-2\alpha-2\beta} L_{,F}], \end{aligned} \quad (10)$$

$$\begin{aligned} \alpha'\gamma' e^{-2\beta} = -\Lambda \\ + \kappa [L(F) + (E^2 e^{-2\alpha-2\beta} - B^2 e^{-2\beta-2\gamma}) L_{,F}], \end{aligned} \quad (11)$$

$$\begin{aligned} e^{-2\beta}(\alpha'\beta' - \alpha'' - \alpha'^2) = \\ \Lambda + \kappa [-L(F) + B^2 e^{-2\beta-2\gamma} L_{,F}] \end{aligned} \quad (12)$$

$$EB L_{,F} = 0, \quad (13)$$

where $\kappa = 8\pi$ and the prime denotes the differentiation d/dr . From the electromagnetic fields equations (5) we arrive at

$$L_{,F} E = C_1 e^{\alpha+\beta-\gamma} \quad (14)$$

and

$$L_{,F} B = C_2 e^{-\alpha+\beta+\gamma}, \quad (15)$$

where C_1 and C_2 are constants of integration.

Notice that from the Einstein equations we conclude that E or B must be zero, i.e. we cannot have any superposition of these nonlinear electric and magnetic fields in any three-dimensional static spacetime, given by diagonal metric (6), just like in the linear case.

On the other hand, from the invariant (8) and the equations (10)–(15) we see that the self-consistent equations are invariant under the duality mapping

$$\begin{aligned} \alpha &\longrightarrow \gamma, \quad \gamma \longrightarrow \alpha \\ E &\longrightarrow iB, \quad B \longrightarrow iE. \end{aligned} \quad (16)$$

Thus we conclude that if we have a nonlinear electric static solution, then one can obtain the nonlinear magnetic one by making the dual mapping (16).

For solving the field equations we must set $E = 0$ (for magnetic solutions) or $B = 0$ (for electric solutions). In

this case the magnetic solutions can be written using the result of Ref. [4], where it was obtained the general solution for the static (2+1)-spacetime for any nonlinear Lagrangian depending on the electric field (i.e. $B = 0$). These electric solutions have the form (6) with structural functions given by

$$\begin{aligned} e^{2\alpha} = e^{-2\beta} &= -M - \Lambda r^2 + 16\pi \int r[L + E^2 L_{,F}]dr, \\ e^{2\gamma} &= r^2 \end{aligned} \quad (17)$$

and, where M is a constant of integration. Then using the dual mapping (16) we obtain the nonlinear magnetic spacetime given by (6), where

$$\begin{aligned} e^{2\alpha} &= r^2, \\ e^{-2\beta} = e^{2\gamma} &= -M - \Lambda r^2 + 16\pi \int r[L - B^2 L_{,F}]dr. \end{aligned} \quad (18)$$

In this case the invariant is $F = B^2/2$. Then from (15) we have that

$$B(r)L_{,F} = \frac{Const}{r} \quad (19)$$

and we can write

$$L_{,r} = \frac{Const}{r} B_{,r}. \quad (20)$$

In order to obtain the Maxwell limit we must set $Const = -q/4\pi$ and from (18) we obtain the solution in terms of the magnetic field $B(r)$:

$$\begin{aligned} e^{2\alpha} &= r^2, \\ e^{-2\beta} = e^{2\gamma} &= -M - \Lambda r^2 - 4q \int \left[r \int \frac{B_{,r}}{r} dr - B \right] dr. \end{aligned} \quad (21)$$

It is easy to check that for Maxwell case, i.e. when $L = -F/4\pi$, we have $B = q/r$ and then from (21) we obtain the HW magnetic solution (3).

A. Some applications

We present now various particular examples of nonlinear magnetic solutions.

One interesting application of our results arises in the Born-Infeld electrodynamics [5]. In this case the Born-Infeld three-dimensional electric spacetime is given by

$$\begin{aligned} e^{2\alpha} = e^{-2\beta} &= -M - (\Lambda - 2b^2)r^2 - 2b^2r\sqrt{r^2 + q_e^2/b^2} \\ &- 2q_e^2 \ln(r + \sqrt{r^2 + q_e^2/b^2}), \quad e^{2\gamma} = r^2, \end{aligned} \quad (22)$$

and the Lagrangian and the electric field are given by

$$L(F) = -\frac{b^2}{4\pi} \left(\sqrt{1 + 2\frac{F}{b^2}} - 1 \right) =$$

$$\begin{aligned} &-\frac{b^2}{4\pi} \left(\frac{r}{\sqrt{r^2 + q_e^2/b^2}} - 1 \right), \\ E(r) &= \frac{q_e}{\sqrt{r^2 + q_e^2/b^2}}, \end{aligned} \quad (23)$$

where b is the Born-Infeld parameter.

Applying the dual mapping in the form $\alpha \rightleftharpoons \gamma$ and $q_e = i q_m$ we obtain for the Born-Infeld magnetic version

$$\begin{aligned} e^{2\alpha} = r^2, \quad e^{-2\beta} = e^{2\gamma} &= -M - (\Lambda - 2b^2)r^2 \\ &- 2b^2r\sqrt{r^2 - q_m^2/b^2} + 2q_m^2 \ln(r + \sqrt{r^2 - q_m^2/b^2}), \end{aligned} \quad (24)$$

and the Lagrangian and the magnetic field are given by

$$\begin{aligned} L(F) &= -\frac{b^2}{4\pi} \left(\sqrt{1 + 2\frac{F}{b^2}} - 1 \right) = \\ &-\frac{b^2}{4\pi} \left(\frac{r}{\sqrt{r^2 - q_m^2/b^2}} - 1 \right), \\ B(r) &= \frac{q_m}{\sqrt{r^2 - q_m^2/b^2}}. \end{aligned} \quad (25)$$

Other interesting example is the magnetic solution obtained from the electric (2+1)-regular black hole solution [4]

$$\begin{aligned} e^{2\alpha} = e^{-2\beta} &= -M - \Lambda r^2 - q_e^2 \ln(r^2 + a^2), \\ e^{2\gamma} &= r^2 \end{aligned} \quad (26)$$

where M , a , q_e and Λ are free parameters. The Lagrangian and the electric field are given by

$$\begin{aligned} L(r) &= \frac{q_e^2}{8\pi} \frac{(r^2 - a^2)}{(r^2 + a^2)^2}, \\ E(r) &= \frac{q_e r^3}{(r^2 + a^2)^2}. \end{aligned} \quad (27)$$

In this case the metric, the electric field and all invariants behave regularly for all values of r and then this solution is curvature regular everywhere. Applying the dual mapping in the form $\alpha \rightleftharpoons \gamma$ and $q_e = i q_m$ we obtain for the magnetic version

$$\begin{aligned} e^{2\alpha} &= r^2, \\ e^{-2\beta} = e^{2\gamma} &= -M - \Lambda r^2 + q_m^2 \ln(r^2 + a^2) \end{aligned} \quad (28)$$

where M , a , q_m and Λ are free parameters. The Lagrangian and the electric field are given by

$$\begin{aligned} L(r) &= -\frac{q_m^2}{8\pi} \frac{(r^2 - a^2)}{(r^2 + a^2)^2}, \\ B(r) &= \frac{q_m r^3}{(r^2 + a^2)^2}. \end{aligned} \quad (29)$$

The last example is the (2+1)-dimensional magnetic solution with a Coulomb-like field. It is obtained from the (2+1)-dimensional black hole with Coulomb-like electric

field. In this case the energy momentum tensor (4) is traceless. The electrostatic spacetime is given by

$$\begin{aligned} e^{2\alpha} = e^{-2\beta} &= -M - \Lambda r^2 + \frac{4q_e^2}{3r}, \\ e^{2\gamma} &= r^2, \end{aligned} \quad (30)$$

where M is a constant of integration related to the mass at infinity [6]. The Lagrangian and the electric field are given by

$$\begin{aligned} L &= \frac{\sqrt{q_e}}{6\pi} |F|^{3/4} = \frac{q_e^2}{6\pi r^3}, \\ E(r) &= \frac{q_e}{r^2}, \end{aligned} \quad (31)$$

where the bars denote absolute value. Applying the dual mapping we obtain

$$\begin{aligned} e^{2\alpha} &= r^2, \\ e^{-2\beta} = e^{2\gamma} &= -M - \Lambda r^2 - \frac{4q_m^2}{3r}. \end{aligned} \quad (32)$$

The Lagrangian and the magnetic field are given by

$$\begin{aligned} L &= -\frac{q_m^2}{6\pi r^3}, \\ B(r) &= \frac{q_m}{r^2}. \end{aligned} \quad (33)$$

Thus, for any three-dimensional static nonlinear electrodynamics whose Lagrangian L depends on a single invariant $F = F_{ab}F^{ab}/4$, and F_{ab} is given by (7), exist a dual mapping (16) between electrostatic and magnetostatic counterparts.

III. STATIONARY EINSTEIN-NONLINEAR MAXWELL THREE-DIMENSIONAL SPACETIMES

Now we shall consider the generalization of these Einstein-nonlinear Maxwell three-dimensional static fields to include the angular momentum J . In this case one requires invariance under spatial rotations and time translations, then the metric can be written in the following general form

$$ds^2 = -(N^0(r))^2 dt^2 + \frac{dR^2}{G^2(r)} + R^2(r)(N^\theta(r) dt + d\theta)^2. \quad (34)$$

One could find any rotating solution by solving the Einstein-nonlinear Maxwell equations using the electromagnetic field tensor (7) and the metric (34). However, in order to do this we shall find the solutions by means of a rotational Lorentz boost. This kind of transformations can generate from seed static spacetimes new metrics because they are not permitted global coordinate transformations. This procedure of generating new solutions

from seed ones has been quite useful in general relativity. A gravitational field may be locally the same but globally distinct due to differences in the topology of their underlying manifolds. Globally stationary but locally static gravitational fields provide a good example of this fact [7].

In view that for the considered Einstein-nonlinear Maxwell problem there are two branches of solutions, electrostatic and magnetostatic one, we must apply the Lorentz boost separately.

A. Stationary electric nonlinear spacetimes

We first shall consider the electrostatic nonlinear spacetimes. In the linear case, i.e. Maxwell electrodynamics, the extension to consider the rotating electrically charged BTZ black hole has been derived in references [8, 9, 10]. For obtaining the stationary electric spacetime we shall follow essentially the procedure of the mentioned above Ref. [10]. The “rotation boost” may be written as the following transformation [10]

$$\begin{aligned} \tilde{t} &= \frac{t - \omega\theta}{\sqrt{1 - \omega^2}}, \\ \tilde{\theta} &= \frac{\theta - \omega t}{\sqrt{1 - \omega^2}}. \end{aligned} \quad (35)$$

Here the $\tilde{t}$ and $\tilde{\varphi}$ are the old time and angular coordinates, respectively, used to write the static form of the metric.

In this case the new metric may be written in the form (34), where $N^0(r) = N(r)G(r)$ and

$$R^2 = \frac{r^2 - \omega^2 f^2}{1 - \omega^2}, \quad (36)$$

$$G^2 = \left(\frac{dR}{dr}\right)^2 f^2, \quad (37)$$

$$N = \frac{r}{R} \left(\frac{dr}{dR}\right) = \frac{d(r^2)}{d(R^2)}, \quad (38)$$

$$N^\theta = \frac{\omega(f^2 - r^2)}{(1 - \omega^2)R^2}. \quad (39)$$

If one starts with the charged nonrotating electrostatic black hole (17), then $f(r) = e^{\alpha(r)}$.

We shall consider asymptotically anti de Sitter solution, i.e. the cosmological constant is negative and is related to the cosmological length l by $\Lambda = -1/l^2$. Now on this length will be set equal to unity. Thus we have

$$f^2(r) = -M + r^2 + 16\pi \int r[L + E^2 L_{,F}]dr. \quad (40)$$

Applying the boost (35) to the solution (17) we obtain

$$R^2 = r^2 + \frac{\omega^2}{1 - \omega^2} \left(M - 16\pi \int r[L + E^2 L_{,F}]dr \right), \quad (41)$$

$$G^2 = \frac{\left(r - \frac{8\omega^2 \pi r (L + E^2 L_{,F})}{1 - \omega^2}\right)^2}{R^2} \times \left(-M + r^2 + 16\pi \int r [L + E^2 L_{,F}] dr\right), \quad (42)$$

$$N = \left(1 - \frac{8\pi\omega^2 (L + E^2 L_{,F})}{(1 - \omega^2)}\right)^{-1} \quad (43)$$

and

$$N^\theta = \frac{\omega (-M + 16\pi \int r [L + E^2 L_{,F}] dr)}{(1 - \omega^2) R^2}. \quad (44)$$

The Maxwell tensor $F_M = E dr \wedge d\tilde{t}$ transforms into

$$F_M = \frac{E}{\sqrt{1 - \omega^2}} dr \wedge (dt - \omega d\theta). \quad (45)$$

Due to this rotation we have now a magnetic component. Taking into account the eq. (38) we have for the pseudoscalar magnetic field, in the gauge (34), the expression

$$B = \frac{E \omega R}{\sqrt{1 - \omega^2} \left(r - \frac{8\pi\omega^2 r (L + E^2 L_{,F})}{(1 - \omega^2)}\right)}. \quad (46)$$

The linear case is obtained when $L(F) = F = -E^2/2$ and all these formulas became the equations (69)-(72) and (75) of the Ref.[10].

Note that if we set $L(F) = 0$ one gets from (40)-(44) the following metric

$$ds^2 = -\left(r^2 - \frac{Mr^2}{r^2(1 - \omega^2) + \omega^2 M}\right) dt^2 + \frac{dr^2}{-M + r^2} + \left(r^2 + \frac{\omega^2 M}{1 - \omega^2}\right) \left(d\theta - \frac{\omega M}{r^2(1 - \omega^2) + \omega^2 M} dt\right)^2, \quad (47)$$

or equivalently

$$ds^2 = -\left(-\frac{M}{1 - \omega^2} + r^2\right) dt^2 + \frac{dr^2}{-M + r^2} - \frac{2\omega M}{1 - \omega^2} dt d\theta + \left(r^2 + \frac{\omega^2 M}{1 - \omega^2}\right) d\theta^2. \quad (48)$$

This is the uncharged rotating BTZ black hole written in an unusual gauge. To write this metric in the usual gauge we apply the following radial coordinate transformation

$$\rho^2 = r^2 + \frac{\omega^2 M}{1 - \omega^2} \quad (49)$$

and we obtain

$$ds^2 = -(-\tilde{M} + \rho^2) dt^2 + \frac{dr^2}{-\tilde{M} + \rho^2 + J^2/(4\rho^2)} - J dt d\theta - \rho^2 d\theta^2 = -\left(-\tilde{M} + \rho^2 + \frac{J^2}{4\rho^2}\right) dt^2 + \left(-\tilde{M} + \rho^2 + \frac{J^2}{4\rho^2}\right)^{-1} dr^2 + \rho^2 \left(\frac{J}{2\rho^2} dt + d\theta\right)^2, \quad (50)$$

where

$$J = \frac{2\omega M}{1 - \omega^2}, \quad \tilde{M} = \frac{M(1 + \omega^2)}{1 - \omega^2}. \quad (51)$$

So, the nonlinear electric rotating solution (40)-(44) reduces to the uncharged rotating BTZ solution when the electric field vanishes.

An interesting example is obtained by application of this procedure to the nonlinear regular black hole (26). In this case the new rotating nonlinear electric spacetime is given by the metric (34), where $N^0 = NG$ and

$$R^2 = \frac{r^2 - \omega^2 (-M + r^2 - q_e^2 \ln(r^2 + a^2))}{1 - \omega^2} \quad (52)$$

$$G^2 = \frac{\left(r - \omega^2 \left(r - \frac{q_e^2 r}{r^2 + a^2}\right)\right)^2}{r^2 - \omega^2 (-M + r^2 - q_e^2 \ln(r^2 + a^2)) (1 - \omega^2)} \times (-M + r^2 - q_e^2 \ln(r^2 + a^2)), \quad (53)$$

$$N = \frac{(r^2 + a^2) (-1 + \omega^2)}{-r^2 - a^2 + \omega^2 r^2 + \omega^2 a^2 - \omega^2 q_e^2}, \quad (54)$$

and

$$N^\theta = \frac{\omega (-M - q_e^2 \ln(r^2 + a^2))}{r^2 - \omega^2 (-M + r^2 - q_e^2 \ln(r^2 + a^2))}. \quad (55)$$

This metric may be written also as

$$ds^2 = -\frac{q_e^2 \ln(r^2 + a^2) + \omega^2 r^2 - r^2 + M}{-1 + \omega^2} dt^2 + \frac{dr^2}{-M + r^2 - q_e^2 \ln(r^2 + a^2)} + \frac{2\omega (-M - q_e^2 \ln(r^2 + a^2))}{1 - \omega^2} dt d\theta + \frac{r^2 - \omega^2 (-M + r^2 - q_e^2 \ln(r^2 + a^2))}{1 - \omega^2} d\theta^2. \quad (56)$$

Other example is the rotating electric Coulomb-like spacetime, which has the following form

$$R^2 = \frac{r^2 - \omega^2 (-M + r^2 + \frac{4q_e^2}{3r})}{(1 - \omega^2)}, \quad (57)$$

$$G^2 = \frac{(-3r^3 + 3\omega^2 r^3 - 2\omega^2 q_e^2)^2 (-3Mr + 3r^3 + 4q_e^2)}{9r^4 (-3r^3 - 3\omega^2 Mr + 3\omega^2 r^3 + 4\omega^2 q_e^2) (-1 + \omega^2)}, \quad (58)$$

$$N = \frac{3r^3 (-1 + \omega^2)}{-3r^3 + 3\omega^2 r^3 - 2\omega^2 q_e^2}, \quad (59)$$

and

$$N^\theta = \frac{(3Mr - 4q_e^2)\omega}{-3r^3 - 3\omega^2 Mr + 3\omega^2 r^3 + 4\omega^2 q_e^2}. \quad (60)$$

This metric may be written also as

$$\begin{aligned} ds^2 = & -\frac{3Mr + 3\omega^2 r^3 - 3r^3 - 4q_e^2}{3r(-1 + \omega^2)} dt^2 \\ & + \frac{dr^2}{-M + r^2 + 4q_e^2/3r} + 2\omega \frac{-M + \frac{4q_e^2}{3r}}{(1 - \omega^2)} dt d\theta \\ & + \frac{r^2 - \omega^2 \left(-M + r^2 + \frac{4q_e^2}{3r}\right)}{(1 - \omega^2)} d\theta^2. \end{aligned} \quad (61)$$

B. Stationary magnetic nonlinear spacetimes

Now we shall consider the magnetostatic nonlinear spacetimes. In order to write the magnetic rotating spacetime in linear electrodynamics the authors of [11] have applied this kind of transformations. However, the properties of this spacetime were not studied. The extension to consider and study the Einstein-Maxwell rotating magnetic solution has been derived in Ref. [12]. The angular momentum also is added to (2) through a rotational Lorentz boost, as we made before for the electric extension. It is interesting to note that the authors of Ref. [12] interpret the static magnetic field (2) as being composed by a system of two symmetric and superposed electric charges. In this case one of the electric charges is at rest and the other is spinning. Thus there is no electric field since the total electric charge is zero and the magnetic field is generated by the angular electric current.

For obtaining the Einstein-nonlinear Maxwell rotating magnetic solution we shall apply the transformation (35) to the magnetostatic solution (18). Then we obtain the metric

$$\begin{aligned} ds^2 = & \frac{r^2 - \omega^2(-M + r^2 + 16\pi \int r(L - B^2 L_F) dr)}{1 - \omega^2} dt^2 \\ & - \frac{dr^2}{-M + r^2 + 16\pi \int r(L - B^2 L_F) dr} \\ & + \frac{2\omega(-M + 16\pi \int r(L - B^2 L_F) dr)}{1 - \omega^2} dt d\theta \\ & - \frac{-M + r^2 + 16\pi \int r(L - B^2 L_F) dr - r^2 \omega^2}{1 - \omega^2} d\theta^2. \end{aligned} \quad (62)$$

The Maxwell tensor $F_M = B dr \wedge d\tilde{\theta}$ transforms into

$$F_M = \frac{B}{\sqrt{1 - \omega^2}} dr \wedge (d\theta - \omega dt). \quad (63)$$

In this case due to the rotation we have now an electric component.

The metric (62) can be rewritten as

$$ds^2 = -R^2(r)(dt + N^\theta(r) d\theta)^2 + \frac{dR^2}{G^2(r)} + (N^0(r))^2 d\theta^2, \quad (64)$$

where $N^0(r) = N(r)G(r)$ and

$$\begin{aligned} R^2 = & r^2 + \frac{\omega^2}{1 - \omega^2} \left(M - 16\pi \int r[L - B^2 L_F] dr \right), \quad (65) \\ G^2 = & \frac{\left(r - \frac{8\omega^2 \pi r(L - B^2 L_F)}{1 - \omega^2} \right)^2}{R^2} \\ & \times \left(-M + r^2 + 16\pi \int r[L - B^2 L_F] dr \right), \quad (66) \end{aligned}$$

$$N = \left(1 - \frac{8\pi\omega^2(L - B^2 L_F)}{(1 - \omega^2)} \right)^{-1}, \quad (67)$$

$$N^\theta = \frac{\omega(-M + 16\pi \int r[L - B^2 L_F] dr)}{(1 - \omega^2)R^2}, \quad (68)$$

and $dr/dR = NR/r$. Considering the eq. (63) we obtain for the electric component

$$E = \frac{B\omega R}{\sqrt{1 - \omega^2} \left(r - \frac{8\pi\omega^2 r(L - B^2 L_F)}{(1 - \omega^2)} \right)}. \quad (69)$$

Comparing the form of the metric (34) with the line element (64) and the expresions (41)-(44) with the eqs. (65)-(68) we find that both solutions are connected by the same duality mapping, which in this case may be written as

$$dt \longrightarrow id\theta, d\theta \longrightarrow idt, E \longrightarrow iB. \quad (70)$$

Thus, for example, the magnetic rotating counterparts of the solutions (28) and (32) may be obtained applying the duality mapping (70) to the eqs. (56) and (61) respectively.

In conclusion these electric and magnetic rotating metrics have both locally static spacetimes which are connected by a duality mapping. Then the rotating spacetimes are also connected by the same duality mapping.

IV. SOME PROPERTIES OF SPINNING NONLINEAR THREE-DIMENSIONAL SPACETIMES

Both the static and rotating solutions are asymptotically anti-de Sitter. Then it is interesting, for example, to calculate the angular momentum and mass of the rotating solutions.

A. Spinning electric solutions

To identify the angular momentum in the metric of the form (34) we can use the quasilocal formalism [13, 14]. In this case the angular momentum $j(r)$ at a radial boundary r is given by [15]

$$j(r) = \frac{GN^{\theta'} R^3}{N^0 R'},$$

where the prime denotes an ordinary derivative with respect to r . Then J is defined as $J = j(\infty)$. In view that in our case $N^0(r) = N(r)G(r)$ we have that

$$j(r) = \frac{N^{\theta'} R^3}{N R'}. \quad (71)$$

For the rotating nonlinear electric spacetime (40)-(44) we have that the angular momentum at a radial boundary r is given by

$$j(r) = \frac{2\omega}{1-\omega^2} \times \left[M + 8\pi r^2(L + E^2 L_F) - 16\pi \int r(L + E^2 L_F) dr \right]. \quad (72)$$

From the last equation we obtain for the uncharged rotating BTZ black hole ($L(F) = 0$) that $j(r) = 2\omega M/(1-\omega^2)$, i.e. the first relation of eq. (51).

In general the behavior of $j(r)$ at infinity depends on the considered electrodynamics. For the linear case $L = -F/(4\pi) = E^2/(8\pi)$. Then we can write $L + E^2 L_F = -q^2/(8\pi r^2)$ and then

$$j(r) = \frac{2\omega}{1-\omega^2} (M - q_e^2 + 2q_e^2 \ln r). \quad (73)$$

At infinity this expression diverges, as it is well known. This intrinsic logarithmic divergence is due to the fact that the electrostatic potential is logarithmic in the linear case.

The behavior of the angular momentum $j(r)$ at a radial boundary r and its behavior at infinity (J) for nonlinear electric rotating black holes generated by the boost (35) from the nonlinear electrostatic spacetimes (22), (26) and (30) are summarized on Table I. The nonlinear Born-Infeld electric black hole (22) and the regular black hole (26) have an angular momentum, which diverges at infinity. More interesting is the black hole with Coulomb-like field (30). At infinity this rotating spacetime has the same angular momentum J of the uncharged rotating BTZ solution.

In addition, we also shall calculate the mass of solutions. One can use the quasilocal mass formula developed in [13, 14] (see also [15]). In this case, from eq. (34), we have that the quasilocal energy $E(r)$ and the quasilocal mass $M(r)$ at a radial boundary r can be shown to be respectively

$$E(r) = 2 \left(f_{ref} \frac{dR_{ref}}{dr} - f(r) \frac{dR}{dr} \right), \quad (74)$$

TABLE I: Expressions for the angular momentum at a radial boundary r , $j(r)$, and angular momentum, J , for the rotating nonlinear electric Born-Infeld, regular and Coulomb-like black holes.

Rotating Spacetimes	$j(r)$	J
from (22)	$\frac{2\omega}{1-\omega^2} \left(M + 2q_e^2 \ln \left(r + \sqrt{r^2 + \frac{q_e^2}{b^2}} \right) \right)$	∞
(56)	$\frac{2\omega}{1-\omega^2} \left(M + q_e^2 \ln(r^2 + a^2) - \frac{q_e^2 r^2}{r^2 + a^2} \right)$	∞
(61)	$\frac{2\omega}{1-\omega^2} \left(M - \frac{2q_e^2}{r} \right)$	$\frac{2\omega}{1-\omega^2} M$

$$M(r) = 2NGE(r) - j(r)N^\theta, \quad (75)$$

where $f^2(r) = (dR/dr)^{-2} G^2$ and f_{ref} and R_{ref} are the background structural functions which determine the zero of energy. The background metric can be obtained simply by setting constant of integration of a particular solution to some special value, determining this way the reference spacetime. We set $q = 0$ and $M = 0$ in our rotating solutions, arriving at the background spacetime, which corresponds to an asymptotic vacuum anti-de Sitter metric. Then $L_0 dR_0/dr = r/l$ (with $l = 1$). The same background was used in other works for the calculation of the quasilocal mass of some spinning solutions (see for example [14, 15]). The quasilocal mass at spatial infinity is defined as

$$\tilde{M} = m(\infty)$$

The mass and angular momentum may be also obtained applying the formalism of Regge and Teitelboim [16]. For example, for the canonical form of the metric (34) the mass is given by (the details can be found in [12])

$$M(r) = \nu [-R_{,r}(f^2 - f_{ref}^2) + (f^2)_{,r}(R - R_{ref}) - 2f^2(R_{,r} - R_{ref,r})], \quad (76)$$

where $f^2 = (dR/dr)^{-2} G^2$, f_{ref}^2 and R_{ref} are structural functions of a background reference spacetime and ν is a constant factor, which come from the fact that, due to the angle deficit, the integration over azimuthal angle θ is between 0 and $2\pi\nu$ (see [12]).

It can be shown that the quasilocal mass at spatial infinity for the charged spinning BTZ solution and the rotating regular black hole is given by

$$\tilde{M} = \frac{1+\omega^2}{1-\omega^2} M - \frac{2\omega^2 q_e^2}{1-\omega^2} + \frac{2q_e^2}{1-\omega^2} \ln r. \quad (77)$$

This coincidence takes place because the electric solutions (1) and (26) have the same behavior at spatial infinity in static and stationary cases.

TABLE II: Expressions of the quasilocal mass at spatial infinity for the rotating nonlinear electric Born-Infeld, regular and Coulomb-like black holes.

Rotating Spacetimes	$\tilde{M}$	Behavior at ∞
from (22)	$\frac{1+\omega^2}{1-\omega^2} (M + q_e^2 \ln r^2 + q_e^2 \ln 4) + q_e^2$	∞
(56)	$\frac{1+\omega^2}{1-\omega^2} M - \frac{2\omega^2 q_e^2}{1-\omega^2} + \frac{2q_e^2}{1-\omega^2} \ln r$	∞
(61)	$\frac{1+\omega^2}{1-\omega^2} M$	$\frac{1+\omega^2}{1-\omega^2} M$

For the rotating electric Coulomb-like black hole the quasilocal mass at spatial infinity does not diverges. In this case we have $\tilde{M} = \frac{1+\omega^2}{1-\omega^2} M$, i.e. the same value of the uncharged rotating BTZ black hole.

The behavior of the quasilocal mass at infinity, $\tilde{M}$, for nonlinear electric rotating black holes generated from (22), (26) and (30) are summarized on Table II.

Thus $\tilde{M}$ diverges logarithmically for the Born-Infeld and regular black holes, similar to the situation in the angular momentum.

B. Spinning magnetic solutions

For the magnetic solution (62) we can write the angular momentum and mass in the form

$$j(r) = \frac{f(N^0 - R^2 N^{\theta^2})^2}{R N^0} \left(\frac{N^{\theta} R^2}{R^2 N^{\theta^2} - N^{0^2}} \right)' \quad (78)$$

and

$$M(r) = \nu \left[-\sqrt{N^{0^2} - R^2 N^{\theta^2}} (f^2 - f_{ref}^2) \right. \\ \left. (f^2)_{,r} \left(\sqrt{N^{0^2} - R^2 N^{\theta^2}} - K_{ref} \right) \right. \\ \left. - 2f^2 \left(\left(\sqrt{N^{0^2} - R^2 N^{\theta^2}} \right)_{,r} - K_{ref,r} \right) \right], \quad (79)$$

respectively, where the structural functions f , R , N^0 and N^{θ} are given by eqs (64)-(68).

The behavior of the angular momentum and mass at a radial boundary r and its behavior at infinity for rotating nonlinear magnetic spacetimes generated from the static solutions (24), (28) and (32) are summarized on the Tables III and IV respectively.

Comparing the expressions for the angular momentum of rotating magnetic solutions (given in table III) with the electric one (given in the table I), we conclude that

TABLE III: Expressions for the angular momentum at a radial boundary r , $j(r)$, and angular momentum, J , for the rotating nonlinear magnetic spacetimes generated from the static solutions (24), (28) and (32).

Rotating Spacetimes generated from	$j(r)$	J
(24)	$\frac{2\omega}{\omega^2-1} \left(M - 2q_m^2 \ln \left(r + \sqrt{r^2 - \frac{q_m^2}{b^2}} \right) \right)$	∞
(28)	$\frac{2\omega}{\omega^2-1} \left(M - q_m^2 \ln(r^2 + a^2) + \frac{q_m^2 r^2}{r^2 + a^2} \right)$	∞
(32)	$\frac{2\omega}{\omega^2-1} \left(M + \frac{2q_m^2}{r} \right)$	$\frac{2\omega}{\omega^2-1} M$

TABLE IV: Expressions of the mass at spatial infinity for the rotating nonlinear magnetic spacetimes generated from the static solutions (24), (28) and (32).

Rotating Spacetimes generated from	$\tilde{M}$	Behavior at ∞
(24)	$\frac{\omega^2+1}{\omega^2-1} (M - q_m^2 \ln r^2 - q_m^2 \ln 4) - q^2$	∞
(28)	$\frac{\omega^2+1}{\omega^2-1} M + \frac{2q_m^2}{\omega^2-1} - \frac{\omega^2+1}{\omega^2-1} q_m^2 \ln r^2$	∞
(32)	$\frac{\omega^2+1}{\omega^2-1} M$	$\frac{\omega^2+1}{\omega^2-1} M$

the angular momentum of the electric rotating spacetimes and the magnetic rotating one are connected by the duality mapping (70).

The divergence of the mass may be handled as follows [10, 12]. We can rewrite, for example, the eq. (77) in the following form

$$\tilde{M} = \frac{1+\omega^2}{1-\omega^2} M - \frac{2\omega^2 q_e^2}{1-\omega^2} + Div_M(r), \quad (80)$$

where $Div_M(r)$ is the logarithmic term, which diverges if $r \rightarrow \infty$. Now, one encloses the system in a boundary of large radius r_0 . Then, one sums and subtracts $Div_M(r_0)$ to (80) so that this mass is now written as

$$\tilde{M} = M(r_0) + (Div_M(r) - Div_M(r_0)), \quad (81)$$

where

$$M(r_0) = M_0 + Div_M(r_0). \quad (82)$$

The second and third terms of (81) vanish when $r \rightarrow r_0$. One might call $M(r_0)$ the energy within the radius r_0 . This energy differs from M_0 by $-Div_M(r_0)$ which may be interpreted as the electromagnetic energy outside r_0 up to an infinite constant which is absorbed in $M(r_0)$. Then from (82) we have that

$$M_0 = M(r_0) - Div_M(r_0). \quad (83)$$

and this sum is independent of r_0 , finite and equal to the total mass.

Thus in practice the treatment of the mass divergence amounts to forgetting about r_0 and takes as zero the asymptotic limit

$$\lim_{r \rightarrow \infty} Div_M(r) = 0$$

in eq. (80).

The divergence on the angular momentum can be treated in a similar way as the mass divergence.

V. ACKNOWLEDGEMENTS

I thank Carol Muñoz for typing this manuscript. This work was supported by CONICYT through Grant FONDECYT N° 1010485 and by Dirección de Promoción y Desarrollo de la Universidad del Bío-Bío.

-
- [1] M. Bañados, C. Teitelboim and J. Zanelli, Phys. Rev. Lett. **69**, 1849 (1992).
 - [2] E. Hirschmann and D. Welch, Phys. Rev. D **53**, 5579 (1996).
 - [3] Cataldo and Salgado, Phys. Rev. D **54**, 2971 (1996).
 - [4] M. Cataldo and A. García, Phys. Rev. D **61**, 084003 (2000).
 - [5] M. Cataldo and A. García, Phys. Lett. B **456**, 28 (1999).
 - [6] M. Cataldo, N. Cruz, S. del Campo and A. García, Phys. Lett. B **484**, 154 (2000).
 - [7] J. Stachel, Phys. Rev. D **26**, 1281 (1982).
 - [8] G. Clément, Class. Quantum Grav. **10**, L54 (1993).
 - [9] G. Clément, Phys. Lett. B **367**, 70 (1996).
 - [10] C. Martínez, C. Teitelboim and J. Zanelli, Phys. Rev. D **61**, 104013 (2000).
 - [11] M. Cataldo and P. Salgado, Phys. Lett. B **448**, 20 (1999).
 - [12] O.J.C. Días and J.P.S. Lemos, JHEP **0201**, 006 (2002).
 - [13] J.D. Brown and J.W. York, Phys. Rev. D **47**, 1407 (1993).
 - [14] J.D. Brown, J. Creighton and R.B. Mann, Phys. Rev. D **50**, 6394 (1994).
 - [15] K.C.K. Chan, Phys. Lett. B **373**, 296 (1996).
 - [16] T. Regge and C. Teitelboim, Ann. Phys. (NY) **88**, 286 (1974).